

(3) Sesbania aegyptiaca Pers. (HB).

tāmbūlī

Piper betle Linn.

tālī

(1) Corypha umbraculifera Linn. (WIRM 2,p.359; KB 4,p.2570-2571; Nadk. 1,nr.684);

(2) Phyllanthus niruri Linn. (Kai.Nigh., oṣadhivarga 247: = tāmalakī; Rā.nigh., ed. I. Tripāṭhī, parpaṭādi 91-92: = bhūmyāmalī = tāmālikā; Dha.nigh. 3,84: = tāmalakī = bhvāmalakī; Rā.nigh. 5,144-146: = bhūmyāmalī = tāmālikā; Pa.ra. 121-122: = bhūmyāmalakī = tāmalakī);

(3) P. fraternus Webster = P. niruri Hook.f., non Linn. (WIRM 8,p.34, where bhūmyāmalī is identified);

(4) P. urinaria Linn. (acc. to identification of bhūmyāmalī: Nadk. 1,nr.1887; Bt.);

(5) = tālamūlī (Pa.ra. 403): Curculigo orchioides Gaertn. (Bt.; HK; Bpn, guṇūcyādivarga 183; Kai.Nigh., oṣadhivarga 1605cd-1606ab: tālamūlikā);

(6) = pāṭalā (Rā.nigh. 9,150-151);

(7) = tālīśapattra (Vaidyakanighaṇṭu acc. to Vśś):

a) Abies webbiana Lindl. (Bpn, karpūrādivarga 214-215; CD, p.109-110; Dha.nigh., ed. Ojhā 2,53; Dha.nigh., ed. P.V. Sharma 2,53; Dgv, nr.300; DGV 2-3,p.233-235; Dutt; HK; Kai.Nigh., oṣadhivarga 1381cd-1382; Nadk. 1,nr.3; DEP 1,p.4-6);

b) A. pindrow Royle (N.C. Shah in JRIM 8,1,1973,p.51);

c) Taxus baccata Linn. (Bpn, karpūrādivarga 214-215; CD, p.109-110; Dgv, nr.299; DGV 2-3,p.234; DWH 3,p.373-377; Nadk. 1,nr.2441; V. 5,p.21-22; DEP 1,p.5-6);

d) Flacourtia jangomas (Lour.) Raeusch. = F. cataphracta Roxb. (CD, p.109-110; KB 1,p.219-220; Raison; DEP 1,p.5);

e) Rhododendron anthopogon D.Don (Bpn, karpūrādivarga 214-215; CD, p.109-110; Dgv, nr.301; DGV 2-3,p.234);

f) R. campanulatum D. Don (Bpn, karpūrādivarga 214-215);

g) R. lepidotum Wall. ex G. Don (Bpn, karpūrādivarga 214-215);

(8) = tuvarikā (Śabdaratnāvalī acc. to Vśś) = ādhakī (Dha.nigh.