

(5) *EUONYMUS TINGENS* WALL. (Wise); some sources (Bpn, *śākavarga* 112-113; Cakra on Ca.Sū. 27, 116cd-117ab) distinguish two kinds of *kaśeru(ka)* : *rājakaśeru(ka)* and *ciṃcoḍakā*; the latter is also called *ciṃcoḍa*, *ciṃcoṭa(ka)* or *cicoḍha*. *rājakaśeru(ka)* :

- 1) *CYPERUS ROTUNDUS* LINN. (MW ; PW);
- 2) *C. SCARIOSUS* R.BR. = *C. PERTENUIS* ROXB. (MW ; PW ; Vśś);
- 3) *SCIRPUS TUBEROSUS* (Avk ; Nadk. 1, nr. 2256);

the Index Kewensis distinguishes two species of this name :

- a) *ELEOCHARIS TUBEROSA* SCHULT. = *S. TUBEROSUS* ROXB.;
- b) *S. MARITIMUS* LINN. = *S. TUBEROSUS* DESF.;

ciṃcoḍakā : 1) *S. ARTICULATUS* LINN. (Avk ; Chopra ; Nadk. 1, nr. 2253);
 2) *C. ESCULENTUS* LINN. (Bpn, p. 684);
 3) [*MARSILEA DENTATA*] (Vśś).

kākaṇantī = *kākaṇantikā*

- (1) *ABRUS PRECATORIUS* LINN.;
- (2) = *muñja* (HK).

kākatinduka

(1) *DIOSPYROS TOMENTOSA* ROXB. (Avk ; Dutt ; HK ; KB 2, p. 1504-1505 ; Nadk. 1, nr. 852 ; PW ; Vśś);

(2) *D. MELANOXYLON* ROXB. (Bpn, p. 531-532 ; Dutt ; HK ; KB 2, p. 1504-1505 ; V 2, p. 110); Kirtikar and Basu consider 1) and 2) as synonyms, but the Index Kewensis and Hooker describe 1) and 2) as separate species ; see Hooker 3, p. 564 ;

(3) *STRYCHNOS NUX-VOMICA* LINN. (Bpn, p. 561-562 ; Dgv, nr. 218 ; KB 3, p. 1645-1647) ;

(4) *CAPPARIS SEPIARIA* LINN. (KB 1, p. 199-200) = *C. INCANESCENS* DC. (KB 1, p. 199-200).

kākādanī

(1) *ABRUS PRECATORIUS* LINN. (KB 1, p. 764-767) ;

- a) the white variety (MW ; PW ; Vśś) ;
- b) the red variety (Vśś) ;

(2) *CAPPARIS SEPIARIA* LINN. (Avk ; Chopra ; Dgm, p. 160 ; Dutt ; DWH 1, p. 131-132 ; HK ; KB 1, p. 199-200 ; MW ; Na 1, nr. 35 ; Nadk. 1, nr. 453) = *C. INCANESCENS* DC. (KB 1, p. 199-200) ;

(3) *C. SPINOSA* LINN. (Avk ; Dgm, p. 157 ; Nadk. 1, nr. 454 ; V 2, p. 65) ;

(4) *ARDISIA HUMILIS* VAHL = *A. SOLANACEA* ROXB. (Vśś) ;