

- b) [C. MEDICA LINN. VAR. MEDICA] (Chopra; KB 1, p. 486-487; Watt CP, p. 325);
- c) [C. MEDICA LINN. VAR. ACIDA] (KB 1, p. 489-490);
- d) [C. ACIDA ROXB. VAR. 7] (Dutt, p. 127; DWH 1, p. 269); see about the varieties of C. MEDICA LINN. : Cooke 1, p. 200-201; Duthie 1, p. 131-132; Hooker 1, p. 514-515; see about the varieties of C. ACIDA ROXB. : Roxburgh, Flora Indica, p. 589.
- (2) PUNICA GRANATUM LINN. (KB 2, p. 1084-1087).

*bṛhatī*

(1) SOLANUM INDICUM LINN. (Ainslie 2, p. 207-208; Bpn, p. 288-289; Dgv, nr. 229; Dutt; DWH 2, p. 555-557; Dy, p. 637-638; Ghosh; Gul; HK; Kav; KB 3, p. 1755-1757; Nadk. 1, nr. 2329; PA 1, 4, p. 78-104; PW; Vśś);

(2) S. XANTHOCARPUM SCHRAD. ET WENDL. (KB 3, p. 1759-1762; PA 1, 4, p. 78-104; Vśś : = *kaṅṭakārī*);

(3) S. MELONGENA LINN. (Bpn, p. 288-289; PA 1, 4, p. 78-104; Vśś : = *vārtākī*; Wise) = S. LONGUM ROXB. (PA 1, 4, p. 78-104) = S. ESCULENTUM DUN. (Vśś);

(4) S. TORVUM SWARTZ (Bpn, p. 288-289 : *śvetabṛhatī*; PA 1, 4, p. 78-104 : used as an adulterant of S. MELONGENA LINN.);

(5) [S. INSANUM WILLD.] (PA 1, 4, p. 78-104) (see Roxb., p. 191) [= S. MELONGENA LINN. VAR. INSANUM PRAIN] (PA 1, 4, p. 78-104);

(6) S. ACULEATISSIMUM JACQ. (PA 1, 4, p. 78-104 : recently introduced, used as a substitute of S. XANTHOCARPUM).

*bhadramustā*

(1) CYPERUS ROTUNDUS LINN. (Dgv, nr. 337; HB; HK; Kav; KB 4, p. 2638-2640; Nadk. 1, nr. 782);

(2) C. SCARIOSUS R.Br. (Bpn, p. 243-244) = C. PERTENUIS ROXB. (Avk; HB, p. 18, footnote 51);

(3) [C. TUBEROSUS ROTH] (Gul); the Index Kewensis enumerates :

a) C. ROTUNDUS LINN. = C. TUBEROSUS ROXB.;

b) C. ESCULENTUS LINN. = C. TUBEROSUS PURSH;

c) C. STOLONIFERUS RETZ. = C. TUBEROSUS BOJ. HORT. MAURIT.;

d) C. TUBEROSUS ROTTB.;

e) MARISCUS MICROCEPHALUS J. ET C. PRESL = C. TUBEROSUS WILLD. EX KUNTH.